



Package leaflet: Information for the user

Fluenz® Tetra nasal spray suspension

Influenza vaccine (live attenuated, nasal)

Read all of this leaflet carefully before the vaccine is given because it contains important information for you or your child.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, nurse or pharmacist.
- This vaccine has been prescribed for you or your child only. Do not pass it on to others.
- If any of the side effects gets serious, talk to your doctor, nurse or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Fluenz Tetra is and what it is used for
2. What you need to know before you are given Fluenz Tetra
3. How Fluenz Tetra is given
4. Possible side effects
5. How to store Fluenz Tetra
6. Contents of the pack and other information

1. What Fluenz Tetra is and what it is used for

Fluenz Tetra is a vaccine to prevent influenza (flu). It is used in children and adolescents 24 months to less than 18 years of age. Fluenz Tetra will help to protect against the four virus strains contained in the vaccine, and other strains closely related to them.

How Fluenz Tetra works

When a person is given the vaccine, the immune system (the body's natural defence system) will produce its own protection against the influenza virus. None of the ingredients in the vaccine can cause the flu.

Fluenz Tetra vaccine viruses are grown in chicken eggs. Each year the vaccine targets four strains of influenza, following the annual recommendations by the World Health Organisation.

2. What you need to know before you are given Fluenz Tetra

✗ You will not be given Fluenz Tetra:

- if you are **allergic** to gentamicin, gelatin or any of the other ingredients of this vaccine (listed in section 6 "Contents of the pack and other information").
- if you have ever had a severe allergic reaction to eggs or egg proteins. For signs of allergic reactions, see section 4 "Possible side effects".
- if you have a **blood disorder** or a **cancer** that **affects the immune system**.
- if you have been **told by your doctor** that you have a **weakened immune system** as a result of a disease, medicine or other treatment.
- if you are **already taking acetylsalicylic acid** (a substance present in many medicines used to relieve pain and lower fever). This is because of the risk of a very rare but serious disease (*Reye's syndrome*).

→ If any of these apply, **tell your doctor, nurse or pharmacist**.

⚠ Warnings and precautions

Talk to your doctor, nurse or pharmacist before vaccination:

- if the **child is less than 24 months of age**. Children less than 24 months of age should not receive this vaccine because of the risk of side effects.
- if you have **severe asthma** or are currently wheezing.
- if you are in **close contact with someone with a severely weakened immune system** (for example, a bone marrow transplant patient needing isolation).

→ If any of these apply, **tell your doctor, nurse or pharmacist before vaccination**. He or she will decide if Fluenz Tetra is suitable for you.

⚠ Other medicines, other vaccines and Fluenz Tetra

Tell your doctor, nurse or pharmacist if the person being vaccinated is taking, has recently taken or might take any other medicines, including medicines that do not require a prescription.

- **Do not give acetylsalicylic acid** (a substance present in many medicines used to relieve pain and lower fever) **to children** for 4 weeks after vaccination with Fluenz Tetra unless your doctor, nurse or pharmacist tells you otherwise. This is because of the risk of *Reye's syndrome*, a very rare but serious disease that can affect the brain and liver.
- **It is recommended that Fluenz Tetra is not given** at the same time as influenza-specific **antiviral medicines**, such as *oseltamivir* and *zanamivir*. This is because the vaccine may work less effectively.

Your doctor, nurse or pharmacist will decide if Fluenz Tetra can be given at the same time as other vaccines.

⚠ Pregnancy and breast-feeding

- If you are **pregnant**, think you may be pregnant, plan to become pregnant soon or are breast-feeding, **tell your doctor, nurse or pharmacist** before receiving this vaccine. Fluenz Tetra is **not recommended** for women who are pregnant or are breast-feeding.

⚠ Driving and using machines

- Fluenz Tetra has no or negligible influence on the ability to drive and use machines.

3. How Fluenz Tetra is given

Fluenz Tetra will be administered under the supervision of a doctor, nurse or pharmacist.

Fluenz Tetra must only be used as a nasal spray.

Fluenz Tetra must not be injected.

Fluenz Tetra will be given as a spray in each nostril. You can breathe normally while you are given Fluenz Tetra. You do not need to actively inhale or sniff.

Dosage

The recommended dose for children and adolescents is 0.2 ml Fluenz Tetra, administered as 0.1 ml in each nostril.

Children who have not previously had an influenza vaccine will receive a second, follow-up dose after an interval of at least 4 weeks. Follow your doctor, nurse or pharmacist's instructions about if and when your child should return for the second dose.

If you have any further questions on this vaccine, ask your doctor, nurse or pharmacist.

turn over 

4. Possible side effects

Like all medicines, this vaccine can cause side effects, although not everybody gets them. In clinical studies with the vaccine, most side effects were mild in nature and short term.

Ask your doctor, nurse or pharmacist if you want more information about possible side effects from Fluenz Tetra.

Some side effects may be serious

Very rare

(may affect up to 1 in 10,000 people):

- severe allergic reaction: signs of a severe allergic reaction may include shortness of breath and swelling of the face or tongue.

- Tell your doctor straight away or seek urgent medical care if you experience any of the effects above.

Other possible side effects of Fluenz Tetra

Very common

(may affect more than 1 in 10 people):

- runny or stuffy nose
- reduced appetite
- weakness

Common

(may affect up to 1 in 10 people):

- fever
- muscle aches
- headache

Uncommon

(may affect up to 1 in 100 people):

- rash
- nose bleed
- allergic reactions

Reporting of side effects

- If you get any side effects, talk to your doctor, nurse or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine.

United Kingdom

Yellow Card Scheme

Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store